

On the Rankin-Selberg problem: higher power moments of the Riesz mean error term

Yoshio Tanigawa(Nagoya), Wenguang Zhai and Deyu Zhang(Jinan)

Science in China Series A **51**(1)(2008), 148-160.

Abstract

Let $\Delta_1(x; \varphi)$ be the error term of the first Riesz means of the Rankin-Selberg zeta function. We study the higher power moments of $\Delta_1(x; \varphi)$ and derive an asymptotic formula for 3-rd, 4-th and 5-th power moments by using Ivić's large value arguments.

1 Introduction and main results

Let $\varphi(z)$ be a holomorphic form of weight κ with respect to the full modular group $SL_2(\mathbb{Z})$, and denote by $a(n)$ the n -th Fourier coefficient of $\varphi(z)$. Suppose that $\varphi(z)$ is normalized such that $a(1) = 1$ and $T(n)\varphi = a(n)\varphi$ for every $n \in \mathbb{N}$, where $T(n)$ is the Hecke operator of order n . Let

$$c_n = n^{1-\kappa} \sum_{m^2|n} m^{2(\kappa-1)} \left| a\left(\frac{n}{m^2}\right) \right|^2.$$

Deligne's [1] proved that $|a(n)| \leq n^{(\kappa-1)/2} d(n)$, which implies $c_n \ll n^\varepsilon$, where $d(n)$ is the Dirichlet divisor function. The classical Rankin-Selberg problem is to study the upper bound of the error term

$$(1.1) \quad \Delta(x; \varphi) := \sum_{n \leq x} c_n - Cx,$$

2000 Mathematics Subject Classification: 11N37, 11F30.

Key Words: The Rankin-Selberg problem, Riesz mean, Voronoï type formula, Power moment.

This work is supported by National Natural Science Foundation of China(Grant No. 10301018) and National Natural Science Foundation of Shandong Province(Grant No. 2006A31).

where C is an explicit constant. Rankin [7] proved that

$$(1.2) \quad \Delta(x; \varphi) = O(x^{\frac{3}{5}}),$$

which was stated by Selberg [8] again without proof. However, no improvement of (1.2) has been obtained after Rankin and Selberg. Ivić [2] proved that $\Delta(x; \varphi) = \Omega_{\pm}(x^{\frac{3}{8}})$ and conjectured that $\Delta(x; \varphi) = O(x^{\frac{3}{8}+\varepsilon})$.

In [5], Ivić, Matsumoto and Tanigawa studied the Riesz mean of the type

$$D_{\rho}(x; \varphi) := \frac{1}{\Gamma(\rho+1)} \sum_{n \leq x} (x-n)^{\rho} c_n.$$

Define the error term $\Delta_{\rho}(x; \varphi)$ by

$$(1.3) \quad D_{\rho}(x; \varphi) = \frac{\pi^2 \kappa R_0}{6\Gamma(\rho+2)} x^{\rho+1} + \frac{Z(0)}{\Gamma(\rho+1)} x^{\rho} + \Delta_{\rho}(x; \varphi),$$

where

$$R_0 = \frac{12(4\pi)^{\kappa-1}}{\Gamma(\kappa+1)} \iint_{\mathfrak{F}} y^{\kappa-2} |\varphi(z)|^2 dx dy,$$

$$Z(s) = \sum_{n=1}^{\infty} c_n n^{-s}, \quad \Re s > 1,$$

the integral being taken over a fundamental domain $\mathfrak{F}$ of $SL_2(\mathbb{Z})$ and $Z(s)$ can be continued to the whole plane.

Ivić, Matsumoto and Tanigawa [5] studied the relation between $\Delta(x; \varphi)$ and $\Delta_1(x; \varphi)$. They proved if $\Delta_1(x; \varphi) = O(x^{\alpha})$ holds for some $\alpha \geq 0$, then $\Delta(x; \varphi) = O(x^{\alpha/2})$. They also proved that

$$(1.4) \quad \Delta_1(x; \varphi) = O(x^{\frac{6}{5}})$$

and

$$(1.5) \quad \int_1^T \Delta_1^2(x; \varphi) dx = \frac{2}{13} (2\pi)^{-4} \left(\sum_{n=1}^{\infty} c_n^2 n^{-7/4} \right) T^{13/4} + O(T^{3+\varepsilon}).$$

Recently, Ivić [3] studied the fourth power moment of $\Delta_1(x; \varphi)$ and proved that

$$(1.6) \quad \int_1^T \Delta_1^4(x; \varphi) dx \ll T^{11/2+\varepsilon}.$$

Note that (1.6) is the best possible up to ε . Both of (1.5) and (1.6) support the following

Conjecture. The pointwise estimate

$$(1.7) \quad \Delta_1(x; \varphi) \ll x^{9/8+\varepsilon}$$

holds.

In this paper we shall prove that for the fourth-power moment of $\Delta_1(x; \varphi)$, we can get an asymptotic formula, which substantially improves Ivić's estimate (1.6). More generally, we shall establish the asymptotic formula for the k -th power moment of $\Delta_1(x; \varphi)$ with $k = 3, 4, 5$. We first prove the following Theorem 1, which is an upper bound result.

Theorem 1. Suppose $A \geq 0$ is a fixed constant, then we have

$$(1.8) \quad \int_T^{2T} |\Delta_1(x; \varphi)|^A dx \ll \begin{cases} T^{1+9A/8} \log^{39} T, & \text{if } 0 \leq A \leq 16/3, \\ T^{(3+6A)/5} \log^5 T, & \text{if } A > 16/3. \end{cases}$$

Remark 1. From Theorem 1 we know that the estimate

$$(1.9) \quad \int_T^{2T} |\Delta_1(x; \varphi)|^{A_0} dx \ll T^{1+9A_0/8+\varepsilon}$$

holds for $A_0 = 16/3$. The value of A_0 for which (1.9) holds is closely related to the upper bound estimate of $\Delta_1(x; \varphi)$. The exponent $16/3$ follows from the estimate (1.4).

Before stating the asymptotic results, we introduce some notations. Suppose $f : \mathbb{N} \rightarrow \mathbb{R}$ is any function such that $f(n) \ll n^\varepsilon$, $k \geq 2$ is a fixed integer. Define

$$(1.10) \quad s_{k;l}(f) := \sum_{\sqrt[4]{n_1} + \dots + \sqrt[4]{n_l} = \sqrt[4]{n_{l+1}} + \dots + \sqrt[4]{n_k}} \frac{f(n_1) \cdots f(n_k)}{(n_1 \cdots n_k)^{7/8}} \quad (1 \leq l < k),$$

$$(1.11) \quad B_k(f) := \sum_{l=1}^{k-1} \binom{k-1}{l} s_{k;l}(f) \cos \frac{\pi(k-2l)}{4}.$$

We shall use $s_{k;l}(f)$ to denote both of the series (1.10) and its value.

Suppose $A_0 > 3$ is a real number, define

$$K_0 := \min\{n \in \mathbb{N} : n \geq A_0, 2|n\}, \quad b(k) := 4^{k-2} + \frac{k-2}{8},$$

$$\sigma(k, A_0) := \frac{3(A_0 - k)}{4(A_0 - 2)}, \quad 3 \leq k < A_0,$$

$$\delta_1(k, A_0) := \frac{\sigma(k, A_0)}{4b(K_0)}, \quad \delta_2(k, A_0) := \frac{\sigma(k, A_0)}{4b(k) + 4\sigma(k, A_0)}.$$

Theorem 2. Let $A_0 > 5$ be a real number such that (1.9) holds, then for any integer $3 \leq k < A_0$, we have the asymptotic formula

$$(1.12) \quad \int_1^T \Delta_1^k(x; \varphi) dx = \frac{B_k(c)}{(8+9k)2^{3k-4}\pi^{2k}} T^{1+9k/8} + O(T^{1+9k/8-\delta_1(k, A_0)+\varepsilon}).$$

Remark 2. From Theorem 1 we see that $k \in \{3, 4, 5\}$, we can get the asymptotic formula (1.12). Moreover, if the estimate $\Delta_1(x; \varphi) \ll x^{19/16-\delta}$ holds for some small $\delta > 0$, then the asymptotic formula (1.12) holds for $k = 6$.

The constant $\delta_1(k, A_0)$ is very small when $k = 3, 4, 5$. In these three cases, we have the following Theorem 3.

Theorem 3. If $k = 3, 4, 5$, then

$$(1.13) \quad \int_1^T \Delta_1^k(x; \varphi) dx = \frac{B_k(c)}{(8+9k)2^{3k-4}\pi^{2k}} T^{1+9k/8} + O(T^{1+9k/8-\delta_2(k, 16/3)+\varepsilon}).$$

Remark 3. Note that

$$\begin{aligned} \delta_2(3, 16/3) &= 7/248 > 1/36, & \delta_2(4, 16/3) &= 3/662 > 1/221, \\ \delta_2(5, 16/3) &= 3/5192 > 1/1731. \end{aligned}$$

So we get the following corollary.

Corollary 1. We have the following asymptotic formulas,

$$\begin{aligned} \int_1^T \Delta_1^3(x; \varphi) dx &= \frac{B_3(c)}{1120\pi^6} T^{35/8} + O(T^{35/8-1/36+\varepsilon}), \\ \int_1^T \Delta_1^4(x; \varphi) dx &= \frac{B_4(c)}{11264\pi^8} T^{11/2} + O(T^{11/2-1/221+\varepsilon}), \\ \int_1^T \Delta_1^5(x; \varphi) dx &= \frac{B_5(c)}{108544\pi^{10}} T^{53/8} + O(T^{53/8-1/1731+\varepsilon}). \end{aligned}$$

2 Preliminary Lemmas

Lemma 2.1. Suppose $x > 1$ is a real number. For any $N \geq 1$, define

$$(2.1) \quad \mathcal{R}_1(x; N) = (2\pi)^{-2} x^{9/8} \sum_{n \leq N} \frac{c_n}{n^{7/8}} \cos\left(8\pi \sqrt[4]{nx} - \frac{\pi}{4}\right).$$

If $1 \ll N \ll x^2$, then

$$(2.2) \quad \Delta_1(x; \varphi) = \mathcal{R}_1(x; N) + O(x^{1+\varepsilon} + x^{3/2+\varepsilon} N^{-1/2}).$$

Proof. (2.2) is a special case of Lemma 2 of [5]. $\square$

Lemma 2.2. *Let S be an inner-product vector space over $\mathbb{C}$, (a, b) denote the inner product in S and $\|a\|^2 = (a, a)$. Suppose that $\xi, \varphi_1, \dots, \varphi_R$ are arbitrary vectors in S . Then*

$$\sum_{l \leq R} |(\xi, \varphi_l)|^2 \leq \|\xi\|^2 \max_{l_1 \leq R} \sum_{l_2 \leq R} |(\varphi_{l_1}, \varphi_{l_2})|.$$

Proof. This is the well-known Halasz-Montgomery inequality. See (A.30) of Ivić [4]. $\square$

Lemma 2.3. *Suppose $k \geq 3$, $(i_1, \dots, i_{k-1}) \in \{0, 1\}^{k-1}$ such that*

$$\sqrt[4]{n_1} + (-1)^{i_1} \sqrt[4]{n_2} + (-1)^{i_2} \sqrt[4]{n_3} + \dots + (-1)^{i_{k-1}} \sqrt[4]{n_k} \neq 0.$$

Then we have

$$|\sqrt[4]{n_1} + (-1)^{i_1} \sqrt[4]{n_2} + (-1)^{i_2} \sqrt[4]{n_3} + \dots + (-1)^{i_{k-1}} \sqrt[4]{n_k}| \gg \max(n_1, \dots, n_k)^{-(4^{k-2}-4^{-1})}.$$

Proof. We follow Tsang's approach [9]. Consider the number field $\mathbb{Q}(\sqrt[4]{n_1}, \dots, \sqrt[4]{n_k})$. Its degree h is one of the following numbers: $1, 4, 4^2, \dots, 4^k$. If $h = 1$, it is trivial. Suppose $h \geq 4$. Let

$$\alpha = \sqrt[4]{n_1} + (-1)^{i_1} \sqrt[4]{n_2} + (-1)^{i_2} \sqrt[4]{n_3} + \dots + (-1)^{i_{k-1}} \sqrt[4]{n_k}$$

and $\alpha = \alpha_1, \alpha_2, \alpha_3, \dots, \alpha_h$ denotes all conjugates of α . Without loss of generality, suppose

$$\begin{aligned} \alpha_{\frac{h}{4}+j} &= -\alpha_j, \quad 1 \leq j \leq h/4; \\ \alpha_{\frac{h}{2}+j} &= \sqrt{-1}\alpha_j, \quad 1 \leq j \leq h/4, \\ \alpha_{\frac{3h}{4}+j} &= -\sqrt{-1}\alpha_j, \quad 1 \leq j \leq h/4. \end{aligned}$$

Since $\alpha_1 \alpha_2 \dots \alpha_h$ is a nonzero integer, we have $|\alpha_1 \alpha_2 \dots \alpha_h| \geq 1$ and hence

$$|\alpha_1 \alpha_2 \dots \alpha_{h/4}| \geq 1,$$

which combining the trivial estimate $|\alpha_j| \leq k \max(n_1, \dots, n_k)$ gives

$$\begin{aligned} |\alpha| &\geq \frac{1}{\prod_{j=2}^{h/4} |\alpha_j|} \geq k^{1-h/4} \max(n_1, \dots, n_k)^{-(h/16-1/4)} \\ &\geq k^{1-4^{k-1}} \max(n_1, \dots, n_k)^{-(4^{k-2}-1/4)} \end{aligned}$$

if noting that $h \leq 4^k$. This completes the proof of Lemma 2.3. $\square$

Lemma 2.4. *Suppose $y > 1$. Define*

$$s_{k;l}(f; y) := \sum_{\substack{\sqrt[4]{n_1} + \dots + \sqrt[4]{n_l} = \sqrt[4]{n_{l+1}} + \dots + \sqrt[4]{n_k} \\ n_j \leq y, 1 \leq j \leq k}} \frac{f(n_1) \cdots f(n_k)}{(n_1 \cdots n_k)^{7/8}} \quad (1 \leq l < k).$$

Then

$$|s_{k;l}(f; y) - s_{k;l}(f)| \ll y^{-3/4+\varepsilon}.$$

Proof. This lemma can be proved by the same argument of Lemma 3.1 of the second author [10], so we omit the details. $\square$

3 Proof of Theorem 1

In order to prove Theorem 1, it suffices for us to study the upper bound of the integral $\int_T^{2T} |\Delta_1(x; \varphi)|^A dx$.

3.1 A large value estimate of $\Delta_1(x; \varphi)$

We first prove the following Theorem 4 via Ivić approach (see, Chapter 13 of [4]).

Theorem 4. *Suppose $T \leq x_1 < x_2 < \dots < x_M \leq 2T$ satisfies $|\Delta_1(x_l; \varphi)| \gg VT$ ($l = 1, 2, \dots, M$) and $|x_j - x_i| \gg V \gg T^{2/21} \mathcal{L}^{34/21}$ ($i \neq j$), then we have*

$$(3.1) \quad M \ll TV^{-3} \mathcal{L}^5 + T^4 V^{-25} \mathcal{L}^{39},$$

where we put $\mathcal{L} = \log T$.

Proof. Suppose $V < T_0$ is a parameter to be determined later. Let I be any subinterval of $[T, 2T]$ of length not exceeding T_0 and let $G = I \cap \{x_1, x_2, \dots, x_M\}$. Without loss of generality, we may suppose $G = \{x_1, x_2, \dots, x_{M_0}\}$.

Suppose $J = \left\lfloor \frac{(1+4\varepsilon)\mathcal{L}-2\log V}{\log 2} \right\rfloor$, then $N = 2^{J+1} \asymp T^{1+4\varepsilon} V^{-2}$. By Lemma 2.1 we get ($T \leq x \leq 2T$)

$$\begin{aligned} \Delta_1(x; \varphi) &\ll T^{\frac{9}{8}} \left| \sum_{n \leq N} \frac{c_n}{n^{7/8}} e(4(nx)^{1/4}) \right| + T^{1-\varepsilon} V \\ &\ll T^{\frac{9}{8}} \left| \sum_{j=0}^J \sum_{n \sim 2^j} \frac{c_n}{n^{7/8}} e(4(nx)^{1/4}) \right| + T^{1-\varepsilon} V. \end{aligned}$$

Squaring, summing over the set G and then using the Cauchy inequality we get that

$$\begin{aligned}
(3.2) \quad \sum_{l \leq M_0} \Delta_1^2(x_l; \varphi) &\ll T^{\frac{9}{4}} \sum_{l \leq M_0} \left| \sum_j \sum_{n \sim 2^j} c_n n^{-7/8} e\left(4(nx_l)^{1/4}\right) \right|^2 \\
&\ll T^{\frac{9}{4}} \mathcal{L} \sum_{l \leq M_0} \sum_j \left| \sum_{n \sim 2^j} c_n n^{-7/8} e\left(4(nx_l)^{1/4}\right) \right|^2 \\
&\ll T^{\frac{9}{4}} \mathcal{L}^2 \sum_{l \leq M_0} \left| \sum_{n \sim 2^{j_0}} c_n n^{-7/8} e\left(4(nx_l)^{1/4}\right) \right|^2
\end{aligned}$$

for some $0 \leq j_0 \leq J$. Let $N_0 = 2^{j_0}$.

Take $\xi = \{\xi_n\}_{n=1}^\infty$ with $\xi_n = c_n n^{-7/8}$ for $n \sim N_0$ and zero otherwise, and take $\varphi_l = \{\varphi_{l,n}\}_{n=1}^\infty$ with $\varphi_{l,n} = e(4(nx_l)^{1/4})$ for $n \sim N_0$ and zero otherwise. Then

$$\begin{aligned}
(\xi, \varphi_l) &= \sum_{n \sim N_0} \frac{c_n}{n^{7/8}} e(4(nx_l)^{1/4}), \\
(\varphi_{l_1}, \varphi_{l_2}) &= \sum_{n \sim N_0} e\left(4n^{1/4}(x_{l_1}^{1/4} - x_{l_2}^{1/4})\right), \\
\|\xi\|^2 &= \sum_{n \sim N_0} \frac{c_n^2}{n^{7/4}} \ll N_0^{-7/4} \sum_{n \sim N_0} c_n^2 \ll N_0^{-3/4} \mathcal{L}^{1+\varepsilon},
\end{aligned}$$

where we used the bound

$$(3.3) \quad \sum_{n \leq x} c_n^2 \ll x \log^{1+\varepsilon} x$$

proved in [6].

From (3.2) and Lemma 2.2 we get

$$\begin{aligned}
(3.4) \quad M_0 V^2 &\ll T^{1/4} \mathcal{L}^2 \sum_{l \leq M_0} \left| \sum_{n \sim N_0} c_n n^{-7/8} e \left(4(n x_l)^{1/4} \right) \right|^2 \\
&\ll \mathcal{L}^{3+\varepsilon} T^{1/4} N_0^{-3/4} \max_{l_1 \leq M_0} \sum_{l_2 \leq M_0} \left| \sum_{n \sim N_0} e \left(4n^{1/4} (x_{l_1}^{1/4} - x_{l_2}^{1/4}) \right) \right| \\
&\ll \mathcal{L}^{3+\varepsilon} T^{1/4} N_0^{1/4} + \frac{\mathcal{L}^{3+\varepsilon} T^{1/4}}{N_0^{3/4}} \max_{l_1 \leq M_0} \sum_{\substack{l_2 \leq M_0, \\ l_2 \neq l_1}} \left| \sum_{n \sim N_0} e \left(4n^{1/4} (x_{l_1}^{1/4} - x_{l_2}^{1/4}) \right) \right|.
\end{aligned}$$

By the Kuzmin-Landau inequality and the exponent pair

$$(1/11, 9/11) = C_{6/11}((0, 1), (1/6, 4/6))$$

we get

$$\begin{aligned}
\sum_{n \sim N_0} e \left(4n^{1/4} (x_{l_1}^{1/4} - x_{l_2}^{1/4}) \right) &\ll \frac{N_0^{3/4}}{|\sqrt[4]{x_{l_1}} - \sqrt[4]{x_{l_2}}|} + \left(\frac{|\sqrt[4]{x_{l_1}} - \sqrt[4]{x_{l_2}}|}{N_0^{3/4}} \right)^{1/11} N_0^{9/11} \\
&\ll \frac{(N_0 T)^{3/4}}{|x_{l_1} - x_{l_2}|} + \left(\frac{|x_{l_1} - x_{l_2}|}{(N_0 T)^{3/4}} \right)^{1/11} N_0^{9/11} \\
&\ll \frac{(N_0 T)^{3/4}}{|x_{l_1} - x_{l_2}|} + T^{-3/44} T_0^{1/11} N_0^{3/4},
\end{aligned}$$

where we used the mean value theorem and the estimate $|x_{l_1} - x_{l_2}| \leq T_0$.

Substituting this estimate into (3.4) we get

$$\begin{aligned}
(3.5) \quad M_0 V^2 &\ll \mathcal{L}^{3+\varepsilon} T^{1/4} N_0^{1/4} + \mathcal{L}^{3+\varepsilon} T^{1/4} N_0^{-3/4} \\
&\quad \times \max_{l_1 \leq M_0} \sum_{\substack{l_2 \leq M_0, \\ l_2 \neq l_1}} \left(\frac{(N_0 T)^{3/4}}{|x_{l_1} - x_{l_2}|} + T^{-3/44} T_0^{1/11} N_0^{3/4} \right) \\
&\ll \mathcal{L}^{3+\varepsilon} T^{1/4} N_0^{1/4} + \mathcal{L}^{4+\varepsilon} T V^{-1} + \mathcal{L}^{3+\varepsilon} M_0 T^{2/11} T_0^{1/11} \\
&\ll \mathcal{L}^{4+\varepsilon} T V^{-1} + \mathcal{L}^{3+\varepsilon} M_0 T^{2/11} T_0^{1/11},
\end{aligned}$$

where we used the facts that $\{x_l\}$ is V -spaced and $N_0 \ll T^{1+4\varepsilon} V^{-2}$. Take $T_0 = V^{22} T^{-2} \mathcal{L}^{-34}$, it is easy to check that $T_0 \gg V$ if $V \gg T^{2/21} \mathcal{L}^{34/21}$. We get for this T_0 that

$$M_0 \ll \mathcal{L}^{4+\varepsilon} T V^{-3}.$$

Now we divide the interval $[T, 2T]$ into $O(1+T/T_0)$ subintervals of length not exceeding T_0 . In each interval of this type, the number of x_l 's is at most $O(\mathcal{L}^{4+\varepsilon}TV^{-3})$. So we have

$$(3.6) \quad M \ll \mathcal{L}^{4+\varepsilon}TV^{-3}\left(1 + \frac{T}{T_0}\right) \ll \mathcal{L}^5TV^{-3} + \mathcal{L}^{39}T^4V^{-25}.$$

This completes the proof of Theorem 4. $\square$

3.2 Proof of Theorem 1

Now we prove Theorem 1. When $A = 0$, Theorem 1 is trivial. When $0 < A < 2$, it follows from (1.6) and the Hölder's inequality. So later we always suppose $A > 2$. It suffices for us to prove the estimate

$$(3.7) \quad \int_T^{2T} |\Delta_1(x; \varphi)|^A dx \ll \begin{cases} T^{1+9A/8} \log^{39} T, & \text{if } 2 < A \leq 16/3, \\ T^{(3+6A)/5} \log^5 T, & \text{if } A > 16/3. \end{cases}$$

Suppose $x^\varepsilon \ll y \leq x/2$. By the definition of $D_\rho(x; \varphi)$ we get

$$\Delta_1(x; \varphi) = \sum_{n \leq x} (x - n)c_n - \frac{\pi^2 \kappa R_0}{12} x^2 - Z(0)x.$$

So we have

$$\begin{aligned} & |\Delta_1(x + y; \varphi) - \Delta_1(x; \varphi)| \\ & \ll \left| \sum_{n \leq x+y} (x + y - n)c_n - \sum_{n \leq x} (x - n)c_n \right| + |(x + y)^2 - x^2| + y \\ & \ll y \left| \sum_{n \leq x} c_n \right| + \left| \sum_{x < n \leq x+y} (x + y - n)c_n \right| + yx \ll yx, \end{aligned}$$

where we used the bound $\sum_{x < n \leq x+y} c_n \ll y$ proved in [5]. So there exists an absolute constant $c_0 > 0$ such that

$$|\Delta_1(x + y; \varphi) - \Delta_1(x; \varphi)| \leq c_0 xy,$$

which implies that if $|\Delta_1(x; \varphi)| \geq 2c_0 xy$, then

$$(3.8) \quad |\Delta_1(x + y; \varphi)| \geq |\Delta_1(x; \varphi)| - |\Delta_1(x + y; \varphi) - \Delta_1(x; \varphi)| \geq c_0 xy.$$

From (3.8) and an argument similar to (13.70) of Ivić [4] we may write

$$(3.9) \quad \int_T^{2T} |\Delta_1(x; \varphi)|^A dx \ll T^{\frac{8+9A}{8}} + \sum_V V \sum_{r \leq N_V} |\Delta_1(x_r; \varphi)|^A,$$

where $T^{1/8} \leq V = 2^m \leq T^{1/5}$, $VT < |\Delta_1(x_r; \varphi)| \leq 2VT$ ($r = 1, \dots, N_V$) and $|x_r - x_s| \geq V$ for $r \neq s \leq N = N_V$.

Suppose first $2 < A < 24$. By Theorem 4 we get

$$\begin{aligned}
(3.10) \quad V \sum_{r \leq N_V} |\Delta_1(x_r; \varphi)|^A &\ll T^A N_V V^{A+1} \\
&\ll \mathcal{L}^5 T^{1+A} V^{A-2} + \mathcal{L}^{39} T^{4+A} V^{A-24} \\
&\ll T^{1+A+\frac{1}{5}(A-2)} \mathcal{L}^5 + T^{1+\frac{9A}{8}} \mathcal{L}^{39}.
\end{aligned}$$

If $A > 24$, then by Theorem 4 again we get

$$\begin{aligned}
(3.11) \quad V \sum_{r \leq N_V} |\Delta_1(x_r; \varphi)|^A &\ll T^A N_V V^{A+1} \\
&\ll \mathcal{L}^5 T^{1+A} V^{A-2} + \mathcal{L}^{39} T^{4+A} V^{A-24} \\
&\ll T^{1+A+\frac{1}{5}(A-2)} \mathcal{L}^5 + T^{\frac{6A-4}{5}} \mathcal{L}^{39} \\
&\ll T^{\frac{6A+3}{5}} \mathcal{L}^5.
\end{aligned}$$

From (3.8)–(3.11) we get (3.7) easily. This completes the proof of Theorem 1.

4 Proofs of Theorem 2 and Theorem 3

Suppose $T \geq 10$ is a real number. It suffices for us to evaluate the integral $\int_T^{2T} \Delta_1^k(x) dx$. Suppose y is a parameter such that $T^\varepsilon < y \leq T^{1/3}$. For any $T \leq x \leq 2T$, define

$$\mathcal{R}_1 = \mathcal{R}_1(x, y) := (2\pi)^{-2} x^{9/8} \sum_{n \leq y} \frac{c_n}{n^{7/8}} \cos(8\pi \sqrt[4]{xn} - \frac{\pi}{4}),$$

$$\mathcal{R}_2 = \mathcal{R}_2(x, y) := \Delta_1(x; \varphi) - \mathcal{R}_1.$$

We shall show that the higher-power moment of $\mathcal{R}_2$ is small and hence the integral $\int_T^{2T} \Delta_1^k(x; \varphi) dx$ can be well approximated by $\int_T^{2T} \mathcal{R}_1^k dx$, which is easy to evaluate.

4.1 Evaluation of the integral $\int_T^{2T} \mathcal{R}_1^k dx$

For simplicity we set $\mathbb{I} = \{0, 1\}$ and

$$\mathbb{N}^k = \{\mathbf{n} : \mathbf{n} = (n_1, \dots, n_k), n_j \in \mathbb{N}, 1 \leq j \leq k\}.$$

For each element $\mathbf{i} = (i_1, \dots, i_{k-1}) \in \mathbb{I}^{k-1}$, put $|\mathbf{i}| = i_1 + \dots + i_{k-1}$. By the elementary formula

$$\cos a_1 \cdots \cos a_k = \frac{1}{2^{k-1}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \cos(a_1 + (-1)^{i_1} a_2 + (-1)^{i_2} a_3 + \dots + (-1)^{i_{k-1}} a_k),$$

we have

$$\begin{aligned} \mathcal{R}_1^k &= (2\pi)^{-2k} x^{\frac{9k}{8}} \sum_{n_1 \leq y} \cdots \sum_{n_k \leq y} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \prod_{j=1}^k \cos(8\pi \sqrt[4]{n_j x} - \frac{\pi}{4}) \\ &= \frac{x^{\frac{9k}{8}}}{(2\pi)^{2k} 2^{k-1}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{n_1 \leq y} \cdots \sum_{n_k \leq y} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \cos(8\pi \sqrt[4]{x} \alpha(\mathbf{n}; \mathbf{i}) - \frac{\pi}{4} \beta(\mathbf{i})), \end{aligned}$$

where

$$\begin{aligned} \alpha(\mathbf{n}; \mathbf{i}) &:= \sqrt[4]{n_1} + (-1)^{i_1} \sqrt[4]{n_2} + (-1)^{i_2} \sqrt[4]{n_3} + \dots + (-1)^{i_{k-1}} \sqrt[4]{n_k}, \\ \beta(\mathbf{i}) &:= 1 + (-1)^{i_1} + (-1)^{i_2} + \dots + (-1)^{i_{k-1}}. \end{aligned}$$

Thus we can write

$$(4.1) \quad \mathcal{R}_1^k = \frac{1}{(2\pi)^{2k} 2^{k-1}} (S_1(x) + S_2(x)),$$

where

$$\begin{aligned} S_1(x) &:= x^{9k/8} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \cos\left(-\frac{\pi \beta(\mathbf{i})}{4}\right) \sum_{\substack{n_j \leq y, 1 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i})=0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}}, \\ S_2(x) &:= x^{9k/8} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{\substack{n_j \leq y, 1 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) \neq 0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \cos\left(8\pi \alpha(\mathbf{n}; \mathbf{i}) \sqrt[4]{x} - \frac{\pi \beta(\mathbf{i})}{4}\right). \end{aligned}$$

First consider the contribution of $S_1(x)$. We have

$$\begin{aligned} (4.2) \quad &\int_T^{2T} S_1(x) dx \\ &= \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \cos\left(-\frac{\pi \beta(\mathbf{i})}{4}\right) \sum_{\substack{n_j \leq y, 1 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i})=0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \int_T^{2T} x^{\frac{9k}{8}} dx. \end{aligned}$$

It is easily seen that if $\alpha(\mathbf{n}; \mathbf{i}) = 0$, then $1 \leq |\mathbf{i}| \leq k-1$. Let $l = |\mathbf{i}|$, then we have

$$\sum_{\substack{n_j \leq y, 1 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i})=0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} = s_{k;l}(c; y),$$

where $s_{k;l}(f; y)$ was defined in Lemma 2.4. By Lemma 2.4 we get

$$(4.3) \quad \int_T^{2T} S_1(x) dx = B_k^*(c) \int_T^{2T} x^{\frac{9k}{8}} dx + O(T^{1+9k/8+\varepsilon} y^{-3/4}),$$

where

$$B_k^*(c) := \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \cos\left(-\frac{\pi\beta(\mathbf{i})}{4}\right) \sum_{\substack{(n_1, \dots, n_k) \in \mathbb{N}^k \\ \alpha(\mathbf{n}; \mathbf{i})=0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}}.$$

For any $\mathbf{i} \in \mathbb{I}^{k-1} \setminus \mathbf{0}$, let

$$S(c; \mathbf{i}) := \sum_{\substack{(n_1, \dots, n_k) \in \mathbb{N}^k \\ \alpha(\mathbf{n}; \mathbf{i})=0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}}.$$

It is easily seen that if $|\mathbf{i}| = |\mathbf{i}'|$ or $|\mathbf{i}| + |\mathbf{i}'| = k$, then

$$S(c; \mathbf{i}) = S(c; \mathbf{i}') = s_{k;|\mathbf{i}|}(c).$$

From $(-1)^j = 1 - 2j$ ($j = 0, 1$) we also have $\beta(\mathbf{i}) = k - 2|\mathbf{i}|$. So we get

$$\begin{aligned} (4.4) \quad B_k^*(c) &= \sum_{l=1}^{k-1} \sum_{|\mathbf{i}|=l} \cos\left(-\frac{\pi\beta(\mathbf{i})}{4}\right) S(c; \mathbf{i}) \\ &= \sum_{l=1}^{k-1} s_{k;l}(c) \cos \frac{\pi(k-2l)}{4} \sum_{|\mathbf{i}|=l} 1 \\ &= \sum_{l=1}^{k-1} \binom{k-1}{l} s_{k;l}(c) \cos \frac{\pi(k-2l)}{4} = B_k(c). \end{aligned}$$

Now we consider the contribution of $S_2(x)$. By the first derivative test

and Lemma 2.3 we get

$$\begin{aligned}
(4.5) \quad \int_T^{2T} S_2(x) dx &\ll T^{\frac{3}{4} + \frac{9k}{8}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{\substack{n_j \leq y, 1 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) \neq 0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8} |\alpha(\mathbf{n}; \mathbf{i})|} \\
&\ll T^{\frac{3}{4} + \frac{9k}{8}} y^{4^{k-2} - 1/4} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{\substack{n_j \leq y, 1 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) \neq 0}} \frac{c_{n_1} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \\
&\ll T^{\frac{3}{4} + \frac{9k}{8}} y^{4^{k-2} - 1/4 + k/8} \ll T^{\frac{3}{4} + \frac{9k}{8}} y^{b(k)}.
\end{aligned}$$

Hence from (4.1)-(4.6) we get

Lemma 4.1. *For any fixed $k \geq 3$, we have*

$$(4.6) \quad \int_T^{2T} \mathcal{R}_1^k dx = \frac{B_k(c)}{(2\pi)^{2k} 2^{k-1}} \int_T^{2T} x^{\frac{9k}{8}} dx + O(T^{1 + \frac{9k}{8} + \varepsilon} y^{-\frac{3}{4}} + T^{\frac{3}{4} + \frac{9k}{8} + \varepsilon} y^{b(k)}).$$

4.2 Higher-power moments of $\mathcal{R}_2$

Taking $N = T$ in the formula (2.2) of Lemma 2.1, we get

$$\begin{aligned}
(4.7) \quad \mathcal{R}_2 &= (2\pi)^{-2} x^{\frac{9}{8}} \sum_{y < n \leq T} \frac{c_n}{n^{7/8}} \cos(8\pi \sqrt[4]{nx} - \pi/4) + O(T^{1+\varepsilon}) \\
&\ll |x^{\frac{9}{8}} \sum_{y < n \leq T} \frac{c_n}{n^{7/8}} e(4\sqrt[4]{nx})| + T^{1+\varepsilon},
\end{aligned}$$

which implies

$$\begin{aligned}
(4.8) \quad \int_T^{2T} \mathcal{R}_2^2 dx &\ll T^{3+\varepsilon} + \int_T^{2T} |x^{\frac{9}{8}} \sum_{y < n \leq T} \frac{c_n}{n^{7/8}} e(4\sqrt[4]{nx})|^2 dx \\
&\ll T^{3+\varepsilon} + T^{13/4} \sum_{y < n \leq T} \frac{c_n^2}{n^{7/4}} + T^3 \sum_{y < m < n \leq T} \frac{c_n c_m}{(mn)^{7/8} (\sqrt[4]{n} - \sqrt[4]{m})} \\
&\ll T^{3+\varepsilon} + \frac{T^{13/4+\varepsilon}}{y^{3/4}} \\
&\ll \frac{T^{13/4+\varepsilon}}{y^{3/4}},
\end{aligned}$$

where we used the estimate (3.3) and the estimate

$$\sum_{y < m < n \leq T} \frac{c_n c_m}{(mn)^{7/8} (\sqrt[4]{n} - \sqrt[4]{m})} \ll T^\varepsilon,$$

which can be proved in a standard way.

Now suppose y satisfies $y^{4b(K_0)} \leq T$. Hence from Lemma 4.1 we get that

$$\int_T^{2T} |\mathcal{R}_1|^{K_0} dx \ll T^{1+9K_0/8+\varepsilon},$$

which implies

$$(4.9) \quad \int_T^{2T} |\mathcal{R}_1|^{A_0} dx \ll T^{1+9A_0/8+\varepsilon}$$

since $A_0 \leq K_0$. From (1.10) and (4.9) we get

$$(4.10) \quad \int_T^{2T} |\mathcal{R}_2|^{A_0} dx \ll \int_T^{2T} (|\Delta_1(x; \varphi)|^{A_0} + |\mathcal{R}_1|^{A_0}) dx \ll T^{1+9A_0/8+\varepsilon}.$$

For any $2 < A < A_0$, from (4.8), (4.10) and Hölder's inequality we get that

$$(4.11) \quad \begin{aligned} \int_T^{2T} |\mathcal{R}_2|^A dx &= \int_T^{2T} |\mathcal{R}_2|^{\frac{2(A_0-A)}{A_0-2} + \frac{A_0(A-2)}{A_0-2}} dx \\ &\ll \left(\int_T^{2T} \mathcal{R}_2^2 dx \right)^{\frac{A_0-A}{A_0-2}} \left(\int_T^{2T} |\mathcal{R}_2|^{A_0} dx \right)^{\frac{A-2}{A_0-2}} \\ &\ll T^{1+\frac{9A}{8}+\varepsilon} y^{-\frac{3(A_0-A)}{4(A_0-2)}}. \end{aligned}$$

Therefore we have

Lemma 4.2. *Suppose $T^\varepsilon \leq y \leq T^{1/4b(K_0)}$, $2 < A < A_0$, then*

$$(4.12) \quad \int_T^{2T} |\mathcal{R}_2|^A dx \ll T^{1+\frac{9A}{8}+\varepsilon} y^{-\frac{3(A_0-A)}{4(A_0-2)}}.$$

4.3 Upper bound of the integral $\int_T^{2T} \mathcal{R}_1^{k-1} \mathcal{R}_2 dx$

In this subsection we shall estimate the integral $\int_T^{2T} \mathcal{R}_1^{k-1} \mathcal{R}_2 dx$. Suppose $T^\varepsilon \leq y \leq T^{1/4b(K_0)}$, which combining Lemma 4.1 implies that

$$\int_T^{2T} |\mathcal{R}_1|^{k-1} dx \ll T^{1+\frac{9(k-1)}{8}}.$$

Thus from (4.7) we get

$$(4.13) \quad \int_T^{2T} \mathcal{R}_1^{k-1} \mathcal{R}_2 dx = \int_T^{2T} \mathcal{R}_1^{k-1} \mathcal{R}_2^* dx + O(T^{1+\frac{9k}{8}-\frac{1}{8}+\varepsilon}),$$

where

$$\mathcal{R}_2^* = (2\pi)^{-2} x^{\frac{9}{8}} \sum_{y < n \leq T} \frac{c_n}{n^{7/8}} \cos(8\pi \sqrt[4]{nx} - \pi/4).$$

Similarly to (4.1) we can write

$$\mathcal{R}_1^{k-1} \mathcal{R}_2^* = \frac{1}{(2\pi)^{2k} 2^{k-1}} (S_3(x) + S_4(x)),$$

where

$$\begin{aligned} S_3(x) &= x^{\frac{9k}{8}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{y < n_1 \leq T} \sum_{\substack{n_j \leq y, 2 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) = 0}} \frac{c_{n_1} c_{n_2} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \cos(-\frac{\pi}{4} \beta(\mathbf{i})), \\ S_4(x) &= x^{\frac{9k}{8}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{y < n_1 \leq T} \sum_{\substack{n_j \leq y, 2 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) \neq 0}} \frac{c_{n_1} c_{n_2} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \cos(8\pi \sqrt[4]{x} \alpha(\mathbf{n}; \mathbf{i}) - \frac{\pi}{4} \beta(\mathbf{i})). \end{aligned}$$

By Lemma 2.4 the contribution of $S_3(x)$ is

$$\begin{aligned} (4.14) \quad \int_T^{2T} S_3(x) dx &\ll \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{y < n_1 \leq T} \sum_{\substack{n_j \leq y, 2 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) = 0}} \frac{c_{n_1} c_{n_2} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8}} \int_T^{2T} x^{\frac{9k}{8}} dx \\ &\ll \sum_{l=1}^{k-1} |s_{k;l}(c; y) - s_{k;l}(c)| \int_T^{2T} x^{\frac{9k}{8}} dx \\ &\ll T^{1+\frac{9k}{8}+\varepsilon} y^{-3/4}. \end{aligned}$$

By the first derivative test and Lemma 2.3 we get

$$\begin{aligned} (4.15) \quad \int_T^{2T} S_4(x) dx &\ll T^{\frac{3}{4}+\frac{9k}{8}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{y < n_1 \leq T} \sum_{\substack{n_j \leq y, 2 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) \neq 0}} \frac{c_{n_1} c_{n_2} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8} |\alpha(\mathbf{n}; \mathbf{i})|} \\ &\ll T^{\frac{3}{4}+\frac{9k}{8}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{y < n_1 \leq k^4 y} \sum_{\substack{n_j \leq y, 2 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) \neq 0}} \frac{c_{n_1} c_{n_2} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8} |\alpha(\mathbf{n}; \mathbf{i})|} \\ &\quad + T^{\frac{3}{4}+\frac{9k}{8}} \sum_{\mathbf{i} \in \mathbb{I}^{k-1}} \sum_{n_1 > k^4 y} \sum_{\substack{n_j \leq y, 2 \leq j \leq k \\ \alpha(\mathbf{n}; \mathbf{i}) \neq 0}} \frac{c_{n_1} c_{n_2} \cdots c_{n_k}}{(n_1 \cdots n_k)^{7/8} n_1^{1/4}} \\ &\ll T^{\frac{3}{4}+\frac{9k}{8}} y^{b(k)}. \end{aligned}$$

From (4.13)–(4.15) we get

$$(4.16) \quad \int_T^{2T} \mathcal{R}_1^{k-1} \mathcal{R}_2 dx \ll T^{1+\frac{9k}{8}+\varepsilon} y^{-3/4} + T^{\frac{3}{4}+\frac{9k}{8}} y^{b(k)} + T^{1+\frac{9k}{8}-\frac{1}{8}+\varepsilon}.$$

4.4 Proof of Theorem 2.

Suppose $3 \leq k < A_0$ and $T^\varepsilon \leq y \leq T^{1/4b(K_0)}$. By the elementary formula $(a+b)^k = a^k + ka^{k-1}b + O(|a^{k-2}b^2| + |b|^k)$, we get

$$(4.17) \quad \int_T^{2T} \Delta_1^k(x; \varphi) dx = \int_T^{2T} \mathcal{R}_1^k dx + k \int_T^{2T} \mathcal{R}_1^{k-1} \mathcal{R}_2 dx \\ + O\left(\int_T^{2T} |\mathcal{R}_1^{k-2} \mathcal{R}_2^2| dx\right) + O\left(\int_T^{2T} |\mathcal{R}_2|^k dx\right).$$

By (4.9), Lemma 4.2 and Hölder's inequality we get

$$(4.18) \quad \int_T^{2T} |\mathcal{R}_1^{k-2} \mathcal{R}_2^2| dx \\ \ll \left(\int_T^{2T} |\mathcal{R}_1|^{A_0} dx\right)^{\frac{k-2}{A_0}} \left(\int_T^{2T} |\mathcal{R}_2|^{\frac{2A_0}{A_0-k+2}} dx\right)^{\frac{A_0-k+2}{A_0}} \\ \ll T^{1+9k/8+\varepsilon} y^{-\frac{3(A_0-k)}{4(A_0-2)}}.$$

Now take $y = T^{1/4b(K_0)}$. From (4.16)–(4.18), Lemma 4.1 and Lemma 4.2 with $A = k$ we get

$$(4.19) \quad \int_T^{2T} \Delta_1^k(x; \varphi) dx = \frac{B_k(c)}{(\sqrt{2}\pi)^k 2^{k-1}} \int_T^{2T} x^{9k/8} dx + O(T^{1+9k/8-\sigma(k, A_0)/4b(K_0)+\varepsilon}) \\ = \frac{B_k(c)}{(\sqrt{2}\pi)^k 2^{k-1}} \int_T^{2T} x^{9k/8} dx + O(T^{1+9k/8-\delta_1(k, A_0)+\varepsilon}).$$

Theorem 2 follows from (4.19) immediately.

4.5 Proof of Theorem 3

Suppose $T^\varepsilon \leq y \leq T^{1/3}$. Using the arguments of Theorem 4 directly to $\mathcal{R}_1$ we get

$$(4.20) \quad \int_T^{2T} |\mathcal{R}_1|^{A_0} dx \ll T^{1+9A_0/8+\varepsilon}$$

holds with $A_0 = 16/3, T^\varepsilon \leq y \leq T^{1/3}$. Thus

$$(4.21) \quad \int_T^{2T} |\mathcal{R}_2|^{A_0} dx \ll T^{1+9A_0/8+\varepsilon}$$

holds with $A_0 = 16/3, T^\varepsilon \leq y \leq T^{1/3}$.

By the same argument as in last subsection, we get that for $T^\varepsilon \leq y \leq T^{1/3}$,

$$(4.22) \quad \int_T^{2T} \Delta_1^k(x; \varphi) dx = \int_T^{2T} \mathcal{R}_1^k dx + O(T^{1+9k/8+\varepsilon} y^{-\sigma(k, 16/3)} + T^{\frac{3}{4} + \frac{9k}{8}} y^{b(k)}).$$

Take $y = T^{1/(4b(k)+4\sigma(k, 16/3))}$. From Lemma 4.1 again we get

$$(4.23) \quad \begin{aligned} \int_T^{2T} \Delta_1^k(x; \varphi) dx &= \frac{B_k(c)}{(2\pi)^k 2^{k-1}} \int_T^{2T} x^{9k/8} dx + O(T^{1+9k/8 - \frac{\sigma(k, 16/3)}{4b(k)+4\sigma(k, 16/3)} + \varepsilon}) \\ &= \frac{B_k(c)}{(2\pi)^k 2^{k-1}} \int_T^{2T} x^{9k/8} dx + O(T^{1+9k/8 - \delta_2(k, 16/3) + \varepsilon}). \end{aligned}$$

Now Theorem 3 follows from (4.23).

References

- [1] P. Deligne, La conjecture de Weil I. Inst. Hautes Etudes Sci. Publ. Math. **43** (1974), 273–307.
- [2] A. Ivić, Large values of certain number-theoretic error terms Acta Arith. **56** (1990), 135–159.
- [3] A. Ivić, On the fourth moment in the Rankin-Selberg problem, submitted.
- [4] A. Ivić, The Riemann zeta-function. John. Wiley and Sons, 1985.
- [5] A. Ivić, K. Matsumoto and Y. Tanigawa, On Riesz means of the coefficients of the Rankin-Selberg series, Math. Proc. Camb. Phil. Soc. **127** (1999), 117–131.
- [6] K. Matsumoto, The mean values and the universality of Rankin-Selberg L-functions. M. Jutila (ed.) et al., Number theory. "Proc. Turku symposium on number theory in memory of K. Inkeri", Turku, Finland, May 31-June 4, 1999. de Gruyter, Berlin, 2001, pp. 201-221.

- [7] R. A. Rankin, Contributions to the theory of Ramanujan's function $\tau(n)$ and similar arithmetical functions II. The order of the Fourier coefficients of integral modular forms, Proc. Cambridge Phil. Soc. **35** (1939), 357–372.
- [8] A. Selberg, Bemerkungen über eine Dirichletsche Reihe, die mit der Theorie der Modulformen nahe verbunden ist, Arch. Math. Naturvid. **43** (1940), 47–50.
- [9] Kai-Man Tsang, Higher-power moments of $\Delta(x)$, $E(t)$ and $P(x)$, Proc. London Math. Soc.(3) **65** (1992), 65–84.
- [10] Wenguang Zhai, On higher-power moments of $\Delta(x)$ (II), Acta Arith. **114** (2004), 35–54.

Yoshio Tanigawa
 Graduate School of Mathematics,
 Nagoya University, Chikusa-ku
 Nagoya 464-8602, Japan
 E-mail: tanigawa@math.nagoya-u.ac.jp

Wenguang Zhai and Deyu Zhang
 School of Mathematical Sciences,
 Shandong Normal University,
 Jian, 250014, Shandong, China
 E-mail: zhaiwg@hotmail.com, zdy-78@yahoo.com.cn